

Important! Read carefully!
Sterile eye drops!

Artelac®

Active ingredient: methylhydroxypropylcellulose

Composition

Active ingredients: 1 ml of solution contains methylhydroxypropylcellulose (HPMC 2906) 3.20 mg, preserved with cetrimide 0.1 mg. The viscosity of the solution is between 7–13 mPa·s.

Other ingredients: sodium monohydrogen phosphate, sodium dihydrogen phosphate, disodium edetate, sorbitol.

Indications

Symptomatic treatment of dehydration manifestations of the cornea and conjunctiva due to impaired lacrimal secretion and functional disorders as a result of topical or systemic diseases or caused by deficient or incomplete eyelid closure.

Contraindications

Hypersensitivity to any of ingredients.

Dosage and administration

Unless otherwise prescribed by doctor, instil 1 drop into the conjunctival sac 3–5 times daily or more frequently if required. The therapy of dry eye requires an individual dosage regimen.

Special remarks

Wearers of soft contact lenses should remove their lenses before instilling Artelac and should wait for at least 15 minutes before inserting them again.

Do not use this medicament beyond the expiry date (refer to outer carton and label).

As in the case of all eye drops, discard any remaining contents 4 weeks after first opening bottle.

Protect the eye drops from light by replacing into outer carton after each usage.

Do not store at temperatures exceeding 25 °C.

Keep all medicines in a safe place out of the reach of children!

Presentation

Dropper bottle of 10 ml solution.

Dr. Gerhard Mann · Chem.-pharm. Fabrik GmbH · 13581 Berlin · Germany

EDV-Nr. 40751

Dr. Mann  Pharma

PB 188/2-E/ROC

愛特淚點眼液

Artelac Eye Drops

衛署藥輸字第 020735 號

成分： 1ml 溶液含：

Hydroxypropylmethylcellulose(HPMC 2906).....	3.20mg
Sorbitol.....	45.00mg
Cetrimide.....	0.1 mg
Disodium phosphate Dodecahydrate.....	q.s.
Sodium Dihydrogen phosphate Dihydrate.....	q.s.
Disodium edetate.....	q.s.
Water for injections.....	q.s.

其溶液黏度為 7-13 mPa.s.

適應症： 眼睛乾澀。

用法用量： 醫師藥師藥劑生指示藥品。

普通每日 3-5 次，若需要可增多，滴 1 滴於眼結膜囊內，治療眼乾症，可視病情調整使用次數。

禁忌症： 對本品所含成分過敏者。

注意： 戴軟性隱形眼鏡於使用本品前，需將隱形眼鏡拿下，用藥後至少停 15 分鐘後再戴上。

過期後不宜使用（失效日期外盒和標籤上有註明）和一般眼藥水一樣，第一次開瓶使用，4 星期後，如未用完，剩下的應拋棄。

避光儲存，每次用完後，應將藥瓶放入盒內。

儲存溫度不可高於 25°C。

應貯存於兒童伸手不及之處。

包裝： 100 公撮以下瓶裝。

製造廠： Dr. Gerhard Mann Chem-pharm. Fabrik GmbH

Brunsbütteler Damm 165-173, 13581 Berlin, Germany

藥商： 武昌貿易有限公司

地址： 台北市南京西路 13 號 8 樓

電話： 2562-6721

EDV-Nr. 40751

PB 188/2-E/ROC