

LIPOSIC EYE GEL

淚保舒眼用凝膠

LIPOSIC EYE GEL

衛署藥輸字第024330號

Composition

1 g contains carbomer 2.0 mg, cetrimide 0.1 mg, medium-chain triglycerides 10.0 mg, sorbitol 48.51 mg.

Other ingredients: sodium hydroxide (for pH adjustment) and purified water.

Properties

Liposic is a three-phase ophthalmic preparation (aqueous, mucous, and lipid phase) intended as artificial lacrymal fluid for the treatment of the dry eye.

Liposic is a thixotropic hydrogel with the viscosity of about 5000mPa.s which is high enough to keep the oil droplets homogeneously dispersed in the gel matrix.

When the gel is instilled into the eye the viscosity of the gel become 10 mPa.s that means it is a solution. The droplets of oil move to the surface of the tear film. When eye blinking the oil forms an even lipid layer to prevent the evaporation of the tear.

Indications

Substitution of tear fluid for management of dry eye conditions such as symptomatic treatment of keratoconjunctivitis sicca.

Dosage and administration

Therapy of dry eye conditions requires an individual dosage regimen. According to the severity of the symptoms, instil one drop into the conjunctival sac 3-5 times daily or more frequently.

Liposic is a liquid gel. It is comfortable to use, when you follow the instructions for instillation. Due to its high viscosity, it is particularly suitable for the treatment of severe dry eye symptoms. Liposic forms a protective film which keeps the surface of the eye moist for a long time. Even in severe cases, only relatively few applications (4-5 per day) should be sufficient to provide relief and bring about improvement.

Use in children: your doctor will advise you of the correct dose.

Contraindication

Hypersensitivity to any component of this product.

Precautions

Contact lenses should be removed prior to administration, and may be inserted again 30 minutes after Liposic has been instilled.

Liposic should only be used in pregnancy and lactation if your doctor thinks it is necessary.

Even when used as indicated, this medicinal product may impair visual acuity for about five minutes, and patients should exercise caution when driving vehicles or operating machinery.

Drug interactions

If other topical ophthalmic agents are to be used along with Liposic, Liposic should always be the last medication to be administered after an interval of approx. 5 minutes.

Side effects

Usually Liposic does not cause any side effects. There is a slight possibility that irritation and itching of the eye may occur in a very small number of people.

Overdose

Liposic is very well tolerated and can be used without restricting the frequency of applications. Care should be taken to administer small drops to avoid sticky eyelids. Any overdose which might occur is, therefore, of no clinical relevance.

Storage

Do not store at temperature exceeding 25 °C.

Do not use this product beyond the expiry date. Discard any remaining contents 4 weeks after opening the tube.

Presentation

Tube of 10 g, in pack sizes of one or three tubes.

Keep all medicines in a safe place out of the reach of children!

Manufacturer:

Dr. Gerhard Mann, Chem.-pharm. Fabrik GmbH · D-13581 Berlin, Germany

【成分】 每公克含：

Carbomer.....	2 mg
Medium chain triglycerides.....	10 mg
Sorbitol.....	48.510 mg
Cetrimide.....	0.100 mg
Sodium hydroxide solution.....	0.832 mg
Purified water.....	q.s.

【劑型】 眼用凝膠劑

【適應症】 乾眼症。

【用法用量】 本藥須由醫師處方使用。

依照病情輕重，白天及睡前約3-5次，若有需要可增加使用次數，每次一滴點於結膜囊內，使用時隱形眼鏡需取下。

【禁忌症】 對本品任何成分過敏者應禁用。
對 Cetrimide 過敏者應禁用。

【注意事項】 使用前應把隱形眼鏡取下，點後隔30分鐘再戴上。
假如醫師認為有需要時方可使用於孕婦和授乳婦。
縱使依指示使用，本品點後會有大約5分鐘的視力模糊現象，故開車和操作機器者宜注意。

【副作用】 本品通常不會造成任何副作用，極少病人偶有眼睛暫時的刺激感和發癢。

【交互作用】 假如與其他點眼劑併用，請至少隔開5分鐘，且本品應最後使用。

【過量】 應避免使用時接觸到眼睛，應避免任何過量使用。

【貯存】 不要存放於溫度25°C以上。
產品逾期不要使用。

和其他眼睛用藥一樣，開啟後一個月未用完應丟棄。
置於兒童伸手不及之處。

【包裝】 100公克以下軟管裝。

製造廠：Dr. Gerhard Mann Chem.-pharm. Fabrik GmbH

廠址：Brunsbütteler Damm 165-173, 13581 Berlin, Germany.

藥商：武昌貿易有限公司

地址：台北市南京西路13號8F